

University of Groningen

Two-Step Macrocycle Synthesis by Classical Ugi Reaction

Abdelraheem, Eman M M; Khaksar, Samad; Kurpiewska, Katarzyna; Kalinowska-Tłuścik, Justyna; Shaabani, Shabnam; Dömling, Alexander

Published in:
The Journal of Organic Chemistry

DOI:
[10.1021/acs.joc.7b02984](https://doi.org/10.1021/acs.joc.7b02984)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Abdelraheem, E. M. M., Khaksar, S., Kurpiewska, K., Kalinowska-Tłuścik, J., Shaabani, S., & Dömling, A. (2018). Two-Step Macrocycle Synthesis by Classical Ugi Reaction. *The Journal of Organic Chemistry*, 83(3), 1441-1447. <https://doi.org/10.1021/acs.joc.7b02984>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Two-Step Macrocycle Synthesis by Classical Ugi Reaction

Eman M. M. Abdelraheem,^{†,‡,§} Samad Khaksar,[†] Katarzyna Kurpiewska,[§] Justyna Kalinowska-Thućik,[§] Shabnam Shaabani,[†] and Alexander Dömling^{*,†,§}


[†]Department of Drug Design, University of Groningen, A. Deusinglaan 1, 9713 AV Groningen, The Netherlands

[‡]Chemistry Department, Faculty of Science, Sohag University, Sohag, 82524, Egypt

[§]Faculty of Chemistry, Jagiellonian University, Gronostajowa 2, 30-387 Krakow, Poland

Supporting Information

ABSTRACT: The direct nonpeptidic macrocycle synthesis of α -isocyano- ω -amines via the classical Ugi four-component reaction (U-4CR) is introduced. Herein an efficient and flexible two-step procedure to complex macrocycles is reported. In the first step, the reaction between unprotected diamines and isocyanocarboxylic acids gives high diversity of unprecedented building blocks in high yield. In the next step, the α -isocyano- ω -amines undergo a U-4CR with high diversity of aldehydes and carboxylic acids in a one-pot procedure. This synthetic approach is short and efficient and leads to a wide range of macrocycles with different ring sizes.



INTRODUCTION

The Ugi four-component reaction (U-4CR) is a widely used multicomponent reaction (MCR) to provide a general route to diverse peptides, macrocycles, and other complex small molecules.^{1,2} This reaction has emerged as a powerful synthetic method for organic and pharmaceutical targets. Among MCRs, isocyanide-based multicomponent reactions (IMCRs) play an important role in pharmaceutical and drug discovery research^{3–7} and provide access to more diverse, complex, and novel scaffolds including small molecules and macrocycles. Macrocycles as intermediates between small molecules and biologics are useful to target flat, large, and featureless protein–protein interfaces.^{8,9} Artificial macrocycles promise to provide better control over synthesizability and over their physico-chemical properties resulting in drug-like properties. However, there are only very few general and short synthetic routes toward macrocycles. Therefore, we report here such a general and short two-step synthesis of macrocycles using the Ugi reaction.

Macrocycles can be synthesized through MCRs by using bifunctional substrates. Failli et al. first used N,C-unprotected tri- and hexapeptides to synthesize bioactive cyclic hexapeptides.¹⁰ Wessjohann et al. used homobifunctional starting materials to synthesize macrocycles using Ugi reactions.¹¹ Yudin et al. introduced formylaziridines as bifunctional Ugi starting materials to synthesize spectacular macrocycles.^{12,13} Recently, Dömling et al. has shown the great impact of the direct use of bifunctional substrates such as α -isocyano- ω -carboxylic acids¹⁴ and α -carboxylic acid- ω -amines¹⁵ in macrocycle synthesis via the Ugi reaction (Figure 1). Of all six possible permutations of bifunctional substrates for macrocyclizations via the Ugi reaction, three have been already

realized, while the last three still deserve validation: α -isocyano- ω -carboxylic acids, α -carboxylic acid- ω -amines, α -isocyano- ω -amines, α -carboxylic acid- ω -aldehydes, α -isocyano- ω -aldehyde, and α -amino- ω -aldehydes. In light of our extended research interest in MCRs and our previous experience in the chemistry of macrocycles, herein we report the use of α -isocyano- ω -amine for the synthesis of macrocycles via the Ugi-macrocyclization reaction.

RESULTS AND DISCUSSION

The first step of our current work is an extension of our recent report on using α -isocyano- ω -amines as building blocks in the cyclization reaction.¹⁶ We started our study by the synthesis of amino isocyanides via coupling of diamines with isocyanide esters under protecting group free conditions. Their synthesis and isolation is demanding due to the highly polar nature of α,ω -amino isocyanides. Therefore, various solvents such as chloroform, dichloromethane (DCM), methanol, water, tetrahydrofuran, ethanol, and trifluoroethanol were tested at room temperature (Table 1). Screening of different solvents revealed that dioxane was the best solvent for this process. Purification was performed by preparative column chromatography on silica (60–200 μ m) using 1:1 dichloromethane:ethyl acetate as eluent A and ammonia in methanol 5% as eluent B in a gradient method. Under the optimized conditions, ten α -isocyano- ω -amines of different lengths were synthesized from commercially available diamines in good purity and yields, each on a gram scale (Scheme 1).

Received: November 30, 2017

Published: January 12, 2018





Figure 1. Six theoretical possibilities for macrocycle synthesis by classical Ugi 4-CR.

In the next step, the macrocyclic ring closure was carried out by an U-4CR under optimized conditions using 1 equiv of an oxo component and an acid (Scheme 2). The optimization was performed by using *N*-(5-aminopentyl)-5-isocyanopentana-mide, paraformaldehyde, and 2-phenylacetic acid as a model reaction. The reaction did not proceed in 1.0 M methanol solution. The same reaction was carried out in different dilutions of methanol, and it was found that a highly diluted 0.01 M equimolar mixture of reactants in methanol gives the 15-membered macrocycle **6a** in good yields (60%). Although trifluoroethanol (65% yield) was slightly superior to MeOH, we chose MeOH for further scope and limitation studies due to the higher price of TFE. Polar aprotic solvents such as THF and CH₃CN gave the product in moderate yields of 30% and 22%, respectively, at room temperature. Next, different Lewis acids such as ZnCl₂ in MeOH and TFE as a solvent were screened. It was found that ZnCl₂ in MeOH affords product in good yield


Table 1. Optimization of Ugi-4CR



entry	solvent (M)	time (h)	catalyst/conditions	yield (%) ^c
1	MeOH (1.0)	12	rt	traces
2	MeOH (0.1)	12	rt	15
3	MeOH (0.01)	12	rt	48
4	TFE (0.01)	12	rt	65
5	CH ₃ CN (0.01)	12	rt	22
6	THF (0.01)	12	rt	30
7 ^b	MeOH (0.01)	12	ZnCl ₂	43
8 ^b	TFE (0.01)	12	ZnCl ₂	25
9	MeOH (0.01)	24	rt	60
10	MeOH (0.01)	12	sonication	20

^aThe reaction was carried out with *N*-(5-aminopentyl)-5-isocyanopentana-mide (1.0 mmol), paraformaldehyde (1.0 mmol), and 2-phenylacetic acid (1.0 mmol). ^b10 mol % catalyst used. ^cYield of isolated product.

Scheme 1. Synthesized α,ω -Amino Isocyanides with Corresponding Yields




(43%). Under sonication conditions, however, the reaction led to low yield of the product (Table 1).

With the optimized reaction conditions in hand, the scope and limitations of the Ugi-macrocyclization reaction were further investigated by synthesizing 15 different macrocycles (12–17 membered ring size) which are shown in Scheme 2. In this reaction, several commercially available carboxylic acids, aliphatic and aromatic aldehydes, and ketones as oxo-components assemble to afford macrocyclic derivatives in good yields of 33–74% after purification by column chromatography. With aliphatic aldehydes, product was obtained in good yields, up to 50%; however, aliphatic carboxylic acids such as isobutyric acid, butyric acid, and pivalic acid resulted in only trace amounts of product.


To investigate potential intramolecular hydrogen bonds of our compounds, a sulfur-containing macrocycle was treated with *m*-chloroperbenzoic acid (*m*CPBA) in DCM to afford sulfoxide and sulfone. As an example, the reaction of macrocycle **6m** with 1 equiv and 4 equiv of *m*CPBA in DCM afforded sulfoxide **7a** and sulfone **7b** in good yields of 65% and

Scheme 2. Synthesized Macrocycles with Corresponding Yields



77%, respectively, after 4 h. As shown in Scheme 3, these sulfoxide and sulfone functional groups are potentially capable

Scheme 3. Selective Oxidative Modifications of a Sulfur-Containing Macrocycle



to form amide–sulfoxide and amide–sulfone intramolecular hydrogen bonds leading to lower energy conformations of the corresponding macrocycles with interlocked structures which could have a significant impact on biological membrane permeability.

X-ray crystal structures of several macrocycles with different sizes and substituents can further provide some first insight into possible solid-state conformations (Figure 2). For instance, compound 6l shows an intramolecular hydrogen bonding.




Figure 2. X-ray crystal structure of some synthesized macrocycles.

Physicochemical properties are of high importance for the development of drug-like compounds. What is the property profile of our macrocycles? To answer this question, we constructed a random virtual 1000 macrocycle library (SI). We calculated some properties of the library related to drug-likeness including molecular weight, lipophilicity, number of hydrogen bond donors and acceptors, number of rotatable bonds, polar surface area, and moment of inertia (Figure 3). Interestingly, analysis of the library shows that 21% obey the Lipinski rule of 5 (RO5). The cLogP vs MW distribution of a considerable fraction of the chemical space is favorable drug-like with an average MW and cLogP of 572 and 4.1, respectively.

Moreover, punctual analysis of 3D modeled representatives and X-ray structures underline the nonflat shapes of the medium sized rings. Overall, a considerable fraction of our macrocyclic space is predicted to have drug-like properties. This is in accordance with the recent proposal that the chemical space from 500 to 1000 Da remains virtually unexplored and represents a vast opportunity for those prepared to venture into new territories of drug discovery.^{17,18}

CONCLUSIONS

A very mild, straightforward, two-step, rapid, and highly diverse macrocycle (12–17 membered) synthesis pathway via MCRs was introduced. In this strategy, macrocyclic ring closure was performed through Ugi-4CR to afford novel complex compounds with potentially biological and pharmaceutical importance. Moreover, our strategy will allow a unique simple route for the synthesis of nonpeptidic macrocycles. Other macrocyclic scaffolds obtained from different combinations of MCRs and their applications as inhibitors for protein–protein interactions are currently being investigated in our laboratory and will be reported shortly.

EXPERIMENTAL SECTION

General Methods. All chemicals were purchased from commercial suppliers and used without any purification unless otherwise noted. Nuclear magnetic resonance spectra were recorded. Chemical shifts for ¹H NMR are reported as δ values, and coupling constants are in hertz (Hz). The following abbreviations are used for spin multiplicity: s = singlet, bs = broad singlet, d = doublet, t = triplet, q = quartet, quin = quintet, dd = doublet of doublets, ddd = doublet of doublet of doublets, m = multiplet. Chemical shifts for ¹³C NMR reported in ppm relative to the solvent peak. Thin layer chromatography was performed on silica gel plates (0.20 mm thick, particle size 25 μ m). Flash chromatography was performed using RediSep R_f normal-phase silica




Figure 3. Some calculated physicochemical properties of the chemical space of macrocycles. A: cLogP over MW scatter plot, B: cLogP over MW box plot, C: Lipinski RO5 radar plot, D: compound distribution based on Lipinski RO5.

flash columns (silica gel 60 Å, 230–400 mesh). Electrospray ionization mass spectra (ESI-MS) were recorded.

Procedure and Analytical Data for Synthesis of α -Isocyano- ω -amine. A round-bottom flask was charged with a magnet stirrer, the diamine (6.0 equiv), and the α -isocyano- ω -methyl ester (5.0 equiv), and 1,4-dioxane (0.1 M) was added. The reaction mixture was stirred at room temperature overnight. The solvent was removed under reduced pressure, and the residue was purified by column chromatography on silica (eluent: 0–100% AB; A 1:1 mixture of EtOAc:DCM, B: methanol, next with C: methanol containing 5% concd aq ammonia, particle size: 40–63 μ m).

***N*-(5-Aminopentyl)-5-isocyanopentanamide **3a**.** The product was obtained as an oil (55%, 0.580 g). ^1H NMR (500 MHz, CDCl_3) δ 6.58 (t, J = 5.8 Hz, 1H), 3.41–3.34 (m, 2H), 3.15 (q, J = 6.7 Hz, 2H), 2.65 (t, J = 7.1 Hz, 2H), 2.16 (t, J = 7.0 Hz, 2H), 1.74–1.61 (m, 4H), 1.49–1.38 (m, 4H), 1.33–1.24 (m, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 172.3, 155.6, 41.4, 39.2, 35.2, 32.1, 29.2, 28.5, 24.0, 22.5. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{11}\text{H}_{22}\text{N}_3\text{O}$ 212.1758; found 212.1757.

***N*-(3-Aminopropyl)-6-isocyanohexanamide **3b**.**¹⁶ The product was obtained as an oil (60%, 0.591 g). ^1H NMR (500 MHz, CDCl_3) δ 6.62 (bs, 1H), 3.42–3.36 (m, 2H), 3.36–3.29 (m, 2H), 2.77 (t, J = 6.4 Hz, 2H), 2.18 (t, J = 7.5 Hz, 2H), 1.72–1.59 (m, 6H), 1.51–

1.42 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ 172.8, 155.6, 50.2, 41.5, 39.8, 37.7, 36.3, 32.2, 28.8, 25.9.

***N*-(4-Aminobutyl)-3-isocyanopropanamide **3c**.**¹⁶ The product was obtained as an oil (60%, 0.464 g). ^1H NMR (500 MHz, CD_3OD) δ 3.78 (t, J = 6.3 Hz, 2H), 3.26 (t, J = 6.5 Hz, 2H), 2.73 (t, J = 6.5 Hz, 2H), 2.66–2.51 (m, 2H), 1.64–1.51 (m, 4H); ^{13}C NMR (126 MHz, CD_3OD) δ 171.6, 156.8, 42.0, 40.3, 39.1, 36.7, 30.3, 27.9.

***N*-(2-((2-Aminoethyl)thio)ethyl)-3-(1H-indol-3-yl)-2-isocyanopropanamide **3d**.**¹⁶ The product was obtained as an oil (49%, 0.774 g). ^1H NMR (500 MHz, CD_3OD) δ 7.58 (d, J = 8.0 Hz, 1H), 7.36 (d, J = 8.0 Hz, 1H), 7.20 (s, 1H), 7.10 (t, J = 7.5 Hz, 1H), 7.04 (t, J = 7.5 Hz, 1H), 4.56 (t, J = 6.6 Hz, 1H), 3.40–3.35 (m, 2H), 3.32 (p, J = 1.6 Hz, 1H), 3.29–3.18 (m, 2H), 2.72 (t, J = 6.6 Hz, 2H), 2.51 (t, J = 6.6 Hz, 2H), 2.32 (t, J = 7.1 Hz, 2H); ^{13}C NMR (126 MHz, methanol- d_4) δ 167.1, 158.4, 136.6, 127.1, 123.9, 121.2, 118.6, 118.0, 111.0, 107.7, 58.3, 40.1, 39.1, 33.7, 29.7, 29.5.

***N*-(6-Aminohexyl)-2-isocyano-3-phenylpropanamide **3e**.**¹⁶ The product was obtained as an oil (56%, 0.764 g). ^1H NMR (500 MHz, CD_3OD) δ 7.42–7.23 (m, 5H), 3.34 (t, J = 1.7 Hz, 1H), 3.28–3.09 (m, 4H), 2.67 (t, J = 7.2 Hz, 2H), 1.57–1.41 (m, 4H), 1.40–1.31 (m, 4H); ^{13}C NMR (126 MHz, CD_3OD) δ 166.3, 158.7, 135.1, 129.1, 128.3, 127.1, 58.3, 40.9, 39.3, 38.9, 31.7, 28.7, 23.3.

N-(5-Aminopentyl)-4-isocyanobutanamide 3f. The product was obtained as an oil (45%, 0.443 g). ^1H NMR (500 MHz, CD_3OD) δ 3.60–3.53 (m, 1H), 3.37–3.32 (m, 1H), 3.23 (t, J = 7.0 Hz, 1H), 2.99–2.90 (m, 2H), 2.39 (t, J = 8.4, 6.4 Hz, 1H), 2.05–1.97 (m, 2H), 1.77–1.67 (m, 2H), 1.65–1.55 (m, 2H), 1.51–1.41 (m, 2H), 1.34 (t, J = 7.1 Hz, 1H), 1.28 (td, J = 7.2, 1.7 Hz, 1H). ^{13}C NMR (126 MHz, CD_3OD) δ 172.9, 154.9, 40.6, 39.3, 38.6, 32.0, 28.5, 27.0, 25.1, 23.4. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{10}\text{H}_{20}\text{N}_3\text{O}$ 198.1601; found 198.1600.

N-(6-Aminohexyl)-3-isocyanopropanamide 3g. The product was obtained as an oil (42%, 0.413 g). ^1H NMR (500 MHz, CD_3OD) δ 3.83–3.72 (m, 1H), 3.38 (s, 2H), 3.36–3.32 (m, 1H), 3.25 (t, J = 7.0 Hz, 1H), 2.67 (t, J = 7.1, 0.9 Hz, 2H), 2.63–2.55 (m, 1H), 1.62–1.47 (m, 4H), 1.40 (m, 4H). ^{13}C NMR (126 MHz, CD_3OD) δ 170.0, 155.2, 41.0, 40.9, 39.0, 37.6, 35.1, 32.2, 32.1, 29.0. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{10}\text{H}_{20}\text{N}_3\text{O}$ 198.1601; found 198.1599.

N-(2-((2-Aminoethyl)thio)ethyl)-3-isocyanopropanamide 3h. The product was obtained as an oil (38%, 0.381 g). ^1H NMR (500 MHz, CDCl_3) δ 6.96 (t, J = 5.7 Hz, 1H), 3.73 (t, J = 6.7, 3.4 Hz, 2H), 3.51–3.45 (m, 2H), 2.92–2.87 (m, 2H), 2.72–2.63 (m, 4H), 2.62–2.55 (m, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 168.5, 156.8, 41.0, 39.0, 37.9, 35.9, 35.7, 31.6. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_8\text{H}_{13}\text{N}_3\text{OS}$ 202.1009; found 202.1008.

N-(2-((2-Aminoethyl)thio)ethyl)-5-isocyanopentanamide 3i. The product was obtained as an oil (57%, 0.652 g). ^1H NMR (500 MHz, CDCl_3) δ 6.52 (t, J = 5.6 Hz, 1H), 3.47–3.36 (m, 4H), 2.88 (t, J = 6.3 Hz, 2H), 2.71–2.58 (m, 4H), 2.23 (t, J = 7.0 Hz, 2H), 1.85–1.67 (m, 4H). ^{13}C NMR (126 MHz, CDCl_3) δ 172.2, 155.9, 41.4, 41.0, 38.7, 35.6, 35.2, 31.6, 28.5, 22.4. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{10}\text{H}_{20}\text{N}_3\text{OS}$ 230.1322; found 230.1320.

N-(6-Aminohexyl)-6-isocyanohexanamide 3j. The product was obtained as an oil (66%, 0.788 g). ^1H NMR (500 MHz, CDCl_3) δ 3.42 (tt, J = 6.6, 1.8 Hz, 1H), 3.22 (q, J = 6.7 Hz, 2H), 2.93–2.80 (m, 4H), 2.23 (t, J = 7.4 Hz, 1H), 1.76–1.58 (m, 6H), 1.57–1.44 (m, 5H), 1.43–1.35 (m, 5H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.0, 155.4, 41.5, 41.1, 40.6, 39.3, 36.2, 31.3, 29.9, 28.8, 26.1, 26.0, 24.8, 24.5. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{13}\text{H}_{26}\text{N}_3\text{O}$ 240.2070; found 240.2069.

Procedure and Analytical Data for the Macrocyclization Reactions. α -Isocyanoo- ω -amine (1.0 mmol) and aldehyde (1.0 mmol) were stirred at room temperature in MeOH (10 mL) for 1 h. The reaction was diluted to (0.01 M, 100 mL), and then the carboxylic acid (1.0 mmol) was added. The mixture was stirred for 24 h. The solvent was removed under reduced pressure, and the residue was purified using flash chromatography (DCM:MeOH (9:1)).

4-(2-Phenylacetyl)-1,4,10-triazacyclopentadecane-2,11-dione 6a. The product was obtained as a white solid (60%, 0.215 g, mp 164–166 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.66 (t, J = 7.0 Hz, 1H), 7.39–7.26 (m, 5H), 6.41 (t, J = 6.3 Hz, 1H), 3.96 (s, 2H), 3.79 (s, 2H), 3.47 (t, J = 6.0 Hz, 2H), 3.28 (q, J = 7.0 Hz, 2H), 3.18 (q, J = 5.6 Hz, 2H), 2.24 (t, J = 7.1 Hz, 2H), 1.67–1.55 (m, 4H), 1.53–1.44 (m, 4H), 1.10 (q, J = 7.9 Hz, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.5, 172.9, 170.4, 134.5, 128.8, 128.8, 127.2, 53.0, 51.8, 40.8, 38.1, 36.8, 35.2, 28.7, 27.9, 27.8, 23.4, 23.1. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{20}\text{H}_{30}\text{N}_3\text{O}_3$ 360.2282; found 360.2281.

4-(2-(4-Bromophenyl)acetyl)-3-isobutyl-1,4,10-triazacyclopentadecane-2,11-dione 6b. The product was obtained as a white solid (51%, 0.251 g, mp 170–172 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.44 (d, J = 8.3 Hz, 2H), 7.09 (d, J = 8.4 Hz, 2H), 6.59 (t, J = 6.1 Hz, 1H), 4.72 (s, 1H), 3.70 (d, J = 3.9 Hz, 2H), 3.55 (s, 1H), 3.44 (s, 1H), 3.39–3.30 (m, 2H), 3.14–2.81 (m, 2H), 2.34–2.13 (m, 2H), 1.88–1.75 (m, 1H), 1.74–1.33 (m, 10H), 1.23 (m, 2H), 0.89 (dd, J = 11.2, 6.6 Hz, 6H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.5, 172.7, 172.1, 133.8, 131.8, 130.8, 121.1, 40.8, 40.7, 38.8, 36.9, 36.4, 35.1, 28.5, 28.0, 24.9, 24.4, 22.8, 22.7, 22.5. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{24}\text{H}_{37}\text{N}_3\text{O}_3\text{Br}$ 494.2013; found 494.2011.

3-(2-(Methylthio)ethyl)-4-(2-phenylacetyl)-1,4,8-triazacyclotetradecane-2,9-dione 6c. The product was obtained as a white solid (50%, 0.209 g, mp 183–185 °C); A mixture of rotamers is observed and the major of the rotamers taken; ^1H NMR (500 MHz, CDCl_3) δ

7.46–7.29 (m, 3H), 7.29–7.20 (m, 2H), 6.10 (s, 1H), 4.85 (t, J = 7.3 Hz, 1H), 3.77 (s, 2H), 3.73–3.54 (m, 2H), 3.53–3.37 (m, 1H), 3.35–3.20 (m, 1H), 3.18–2.96 (m, 2H), 2.57–2.34 (m, 3H), 2.34–2.21 (m, 2H), 2.12 (s, 3H), 2.08–1.92 (m, 2H), 1.86–1.58 (m, 3H), 1.45 (t, J = 11.1 Hz, 2H), 1.36–1.13 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ 173.6, 172.5, 170.7, 135.0, 129.3, 129.2, 127.5, 59.7, 53.4, 46.5, 41.9, 38.5, 37.8, 36.5, 31.2, 30.2, 29.4, 28.9, 24.2, 15.8. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{22}\text{H}_{34}\text{N}_3\text{O}_3\text{S}$ 420.2315; found 420.2313.

3-Isobutyl-4-(2-phenylacetyl)-1,4,8-triazacyclotetradecane-2,9-dione 6d. The product was obtained as brown oil (74%, 0.296 g); ^1H NMR (500 MHz, CDCl_3) δ 7.42–7.37 (m, 2H), 7.36–7.31 (m, 3H), 6.32 (dd, J = 15.3, 8.5 Hz, 1H), 5.69 (t, J = 6.1 Hz, 1H), 5.24–5.17 (m, 1H), 3.73 (s, 2H), 3.33–3.24 (m, 4H), 3.13–3.02 (m, 2H), 2.18 (t, J = 7.5, 1.9 Hz, 2H), 1.71 (dd, J = 7.2, 5.8 Hz, 2H), 1.66–1.59 (m, 5H), 1.35–1.27 (m, 2H), 1.27–1.19 (m, 2H), 0.92 (dd, J = 8.2, 6.3 Hz, 6H); ^{13}C NMR (126 MHz, CDCl_3) δ 173.6, 170.9, 170.3, 133.6, 129.2, 128.9, 127.6, 73.0, 41.6, 40.8, 38.8, 36.4, 36.0, 35.8, 29.7, 28.9, 26.2, 25.1, 24.5, 23.1, 21.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{36}\text{N}_3\text{O}_3$ 402.2751; found 402.2750.

2-(tert-Butyl)-1-(2-(4-nitrophenyl)acetyl)-1,4,8-triazacyclododecane-3,7-dione 6e. The product was obtained as a white solid (36%, 0.150 g, mp 188–190 °C); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.29 (d, J = 8.2 Hz, 1H), 8.21–8.16 (m, 2H), 7.56–7.52 (m, 2H), 4.67 (s, 1H), 4.21–3.97 (m, 2H), 3.85–3.67 (m, 2H), 3.56 (dd, J = 15.2, 8.4 Hz, 1H), 3.18–2.93 (m, 2H), 2.85 (d, J = 13.4 Hz, 1H), 2.38–2.26 (m, 1H), 2.26–2.11 (m, 1H), 1.80–1.64 (m, 1H), 1.50–1.35 (m, 1H), 1.35–1.17 (m, 1H), 1.12 (d, J = 18.4 Hz, 1H), 1.10–1.02 (m, 1H), 0.89 (s, 9H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ 171.2, 171.0, 169.9, 146.7, 145.2, 131.5, 123.6, 37.9, 37.4, 37.3, 37.0, 36.9, 35.9, 28.2, 27.8, 27.1, 26.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{21}\text{H}_{31}\text{N}_4\text{O}_5$ 419.2289; found 419.2287.

2-(4-Chlorophenyl)-1-(2-phenylacetyl)-1,4,8-triazacyclododecane-3,7-dione 6f. The product was obtained as a yellow oil (42%, 0.179 g); ^1H NMR (500 MHz, CDCl_3) δ 8.66 (t, J = 5.2 Hz, 1H), 7.39–7.31 (m, 3H), 7.31–7.25 (m, 2H), 7.24–7.17 (m, 2H), 7.17–7.04 (m, 1H), 6.37 (t, J = 6.6 Hz, 1H), 3.77 (s, 2H), 3.55–3.39 (m, 4H), 3.28–3.07 (m, 2H), 2.69–2.52 (m, 2H), 1.47 (q, J = 3.6 Hz, 4H). ^{13}C NMR (126 MHz, CDCl_3) δ 177.6, 174.9, 171.9, 134.9, 129.3, 128.8, 128.7, 128.5, 127.0, 63.5, 45.9, 44.2, 38.8, 38.1, 35.4, 26.4, 26.1, 25.3. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}_3\text{Cl}$ 428.1735; found 428.1734.

14-((1H-Indol-3-yl)methyl)-6-(2-phenylacetyl)-9-thia-6,12,15-triazaspiro[4.11]hexadecane-13,16-dione 6g. The product was obtained as a yellow solid (53%, 0.274 g, mp 180–182 °C); ^1H NMR (500 MHz, CDCl_3) δ 8.84 (d, J = 7.3 Hz, 1H), 8.18 (s, 1H), 7.66 (d, J = 7.9 Hz, 1H), 7.36 (s, 1H), 7.34 (d, J = 7.4 Hz, 2H), 7.29 (d, J = 6.6 Hz, 1H), 7.18 (t, J = 7.4 Hz, 3H), 7.15–7.10 (m, 2H), 6.79 (t, J = 2.0 Hz, 1H), 4.67–4.55 (m, 1H), 3.88 (s, 2H), 3.68 (s, 1H), 3.61–3.56 (m, 2H), 3.45 (dd, J = 15.0, 5.4 Hz, 1H), 3.24 (dd, J = 15.0, 8.9 Hz, 1H), 3.06–2.98 (m, 2H), 2.90–2.79 (m, 1H), 2.74–2.61 (m, 1H), 2.56–2.45 (m, 1H), 2.32–2.18 (m, 1H), 1.87–1.77 (m, 1H), 1.58–1.47 (m, 4H), 1.34–1.26 (m, 1H), 1.25–1.15 (m, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 175.4, 173.8, 171.8, 136.3, 134.9, 129.4, 128.8, 128.7, 128.6, 127.3, 127.0, 123.2, 122.0, 119.4, 118.9, 111.1, 72.2, 54.9, 45.7, 43.7, 37.6, 37.2, 36.0, 35.5, 34.7, 26.2, 21.6, 21.0. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{29}\text{H}_{35}\text{N}_4\text{O}_3\text{S}$ 519.2424; found 519.2424.

3-Benzyl-6-(tert-butyl)-7-(2-phenylacetyl)-1,4,7-triazacyclotridecane-2,5-dione 6h. The product was obtained as a white oil (33%, 0.157 g); ^1H NMR (500 MHz, CDCl_3) δ 7.45–7.33 (m, 5H), 7.34–7.21 (m, 3H), 4.75 (s, 1H), 3.99 (s, 2H), 3.68 (d, J = 5.2 Hz, 4H), 3.38 (s, 1H), 3.20–3.08 (m, 2H), 1.63–1.49 (m, 3H), 1.46–1.39 (m, 2H), 1.28 (d, J = 2.9 Hz, 1H), 1.24–1.20 (m, 3H), 1.15–1.09 (m, 4H), 1.07 (s, 9H). ^{13}C NMR (126 MHz, CDCl_3) δ 190.3, 183.6, 134.0, 129.0, 128.7, 127.6, 70.2, 52.0, 47.0, 40.7, 36.4, 35.7, 29.8, 28.6, 26.5, 26.4, 26.2, 26.2. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{29}\text{H}_{40}\text{N}_3\text{O}_3$ 478.2912; found 478.2912.

2-Isobutyl-1-(2-phenylacetyl)-1,4,9-triazacyclotetradecane-3,8-dione 6i. The product was obtained as a yellow oil (45%, 0.180 g); ^1H NMR (500 MHz, CDCl_3) δ 7.39–7.32 (m, 5H), 7.18 (d, J = 5.9 Hz,

1H), 5.56 (t, $J = 6.4$ Hz, 1H), 4.26 (s, 1H), 3.83 (d, $J = 5.9$ Hz, 2H), 3.62–3.54 (m, 2H), 3.32 (q, $J = 5.3$ Hz, 2H), 3.18–3.11 (m, 1H), 3.07–2.99 (m, 1H), 2.43–2.34 (m, 1H), 2.26–2.19 (m, 1H), 2.15–2.08 (m, 1H), 1.87 (t, $J = 7.2$ Hz, 2H), 1.81–1.74 (m, 1H), 1.53–1.48 (m, 2H), 1.46–1.42 (m, 1H), 1.37–1.32 (m, 1H), 1.29–1.26 (m, 1H), 1.23 (d, $J = 6.0$ Hz, 1H), 1.01 (t, $J = 6.5$ Hz, 1H), 0.90 (d, $J = 6.2$ Hz, 6H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.2, 172.8, 172.4, 134.9, 128.9, 128.7, 127.0, 61.1, 48.9, 41.5, 40.7, 37.5, 37.2, 35.2, 29.1, 28.6, 25.0, 23.8, 23.1, 23.0, 22.1. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{23}\text{H}_{36}\text{N}_3\text{O}_3$ 402.2751; found 402.2750.

2-(4-Isopropylphenyl)-1-(2-phenylacetyl)-1,4,8-triazacyclotetradecane-3,7-dione 6j. The product was obtained as a white solid (39%, 0.180 g, mp 166–168 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.40–7.29 (m, 3H), 7.28 (s, 1H), 7.24–7.12 (m, 5H), 6.65 (s, 1H), 5.60 (s, 1H), 3.93–3.60 (m, 4H), 3.62–3.39 (m, 3H), 3.28 (d, $J = 51.2$ Hz, 2H), 3.04–2.80 (m, 1H), 2.73–2.35 (m, 2H), 1.77–1.48 (m, 4H), 1.41 (s, 2H), 1.25 (d, $J = 7.1$ Hz, 6H), 1.23 (s, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 172.8, 171.8, 171.0, 129.8, 129.2, 129.1, 128.9, 128.1, 127.3, 127.1, 126.9, 65.7, 49.8, 42.1, 39.8, 36.8, 36.0, 34.1, 29.0, 26.8, 26.5, 25.6, 24.2. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{28}\text{H}_{38}\text{N}_3\text{O}_3$ 464.2908; found 464.2906.

1-(3-Phenylpropanoyl)-1,4,8-triazacyclotetradecane-3,7-dione 6k. The product was obtained as a white solid (44%, 0.157 g, mp 177–179 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.34 (t, $J = 7.5$ Hz, 2H), 7.25 (q, $J = 9.3$, 7.9 Hz, 3H), 6.39 (t, $J = 5.9$ Hz, 1H), 4.01 (s, 2H), 3.54 (q, $J = 5.9$ Hz, 2H), 3.39–3.29 (m, 4H), 3.02 (t, $J = 7.8$ Hz, 2H), 2.71 (t, $J = 7.9$ Hz, 2H), 2.50 (t, $J = 5.8$ Hz, 2H), 1.63–1.51 (m, 4H), 1.44–1.36 (m, 2H), 1.28–1.20 (m, 2H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.8, 170.9, 170.6, 141.0, 128.6, 128.5, 126.3, 52.3, 50.6, 39.3, 36.3, 35.6, 35.3, 31.2, 28.4, 26.7, 26.3, 25.5. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{20}\text{H}_{30}\text{N}_3\text{O}_3$ 360.2282; found 360.2279.

4-(2-Phenylacetyl)-1-thia-4,7,11-triazacyclotridecane-6,10-dione 6l. The product was obtained as a white solid (62%, 0.216 g, mp 173–174 °C); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.53–8.43 (m, 1H), 8.42 (d, $J = 6.1$ Hz, 1H), 7.35–7.26 (m, 2H), 7.27–7.18 (m, 3H), 3.98 (s, 2H), 3.81 (s, 2H), 3.36–3.25 (m, 6H), 2.69 (m, 2H), 2.63–2.58 (m, 2H), 2.41 (t, $J = 5.8$ Hz, 2H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ 171.6, 170.9, 169.7, 136.5, 129.8, 128.6, 126.7, 52.4, 48.6, 42.6, 40.6, 36.2, 34.5, 30.8, 29.8. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{17}\text{H}_{24}\text{N}_3\text{O}_3\text{S}$ 350.1533; found 350.1532.

4-(2-(4-Bromophenyl)acetyl)-1-thia-4,7,13-triazacyclopentadecane-6,12-dione 6m. The product was obtained as a white solid (58%, 0.264 g, mp 196–198 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.48–7.44 (m, 2H), 7.44–7.40 (m, 1H), 7.19–7.13 (m, 2H), 7.11 (d, $J = 8.2$ Hz, 2H), 3.98 (s, 2H), 3.77 (s, 2H), 3.67 (t, $J = 6.2$ Hz, 2H), 3.57 (d, $J = 3.4$ Hz, 1H), 3.46–3.37 (m, 2H), 3.33–3.22 (m, 2H), 2.90 (t, $J = 6.2$ Hz, 2H), 2.83–2.75 (m, 2H), 2.29 (t, $J = 6.3$ Hz, 2H), 1.67–1.57 (m, $J = 5.7$, 4.8 Hz, 4H). ^{13}C NMR (126 MHz, CDCl_3) δ 174.3, 172.3, 170.4, 133.3, 131.9, 130.8, 121.3, 53.1, 50.1, 40.0, 37.9, 36.7, 34.4, 31.4, 30.4, 27.8, 22.5. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_3\text{SBr}$ 456.0949; found 456.0949.

6-(2-(p-Tolyl)acetyl)-9-thia-6,12,18-triazaspiro[4.14]nonadecane-13,19-dione 6n. The product was obtained as a yellow solid (69%, 0.307 g, mp 195–197 °C); ^1H NMR (500 MHz, CDCl_3) δ 7.53 (t, $J = 8.8$ Hz, 2H), 7.27 (dd, $J = 7.9$, 5.1 Hz, 2H), 7.20 (d, $J = 8.2$ Hz, 1H), 6.25 (s, 1H), 5.49 (t, $J = 6.2$ Hz, 1H), 4.96 (s, 1H), 3.85–3.77 (m, 1H), 3.73 (d, $J = 15.1$ Hz, 1H), 3.54–3.46 (m, 1H), 3.42–3.34 (m, 1H), 3.22–3.10 (m, 1H), 3.08–2.90 (m, 2H), 2.66–2.53 (m, 1H), 2.51–2.41 (m, 2H), 2.41–2.27 (m, 1H), 2.10 (s, 3H), 2.02–1.89 (m,

1H), 1.84–1.71 (m, 3H), 1.60–1.43 (m, 7H), 1.41–1.23 (m, 7H). ^{13}C NMR (126 MHz, CDCl_3) δ 173.0, 172.6, 171.1, 133.8, 131.9, 131.0, 121.1, 59.6, 46.5, 40.3, 39.4, 37.7, 36.2, 30.8, 29.7, 29.7, 28.8, 27.8, 26.7, 25.6, 25.0, 24.2, 15.4. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{25}\text{H}_{39}\text{N}_3\text{O}_3\text{SBr}$ 540.1890; found 540.1890.

General Procedure and Analytical Data for the Synthesis of Sulfoxide Macrocycle. Macrocycle **6q** (1.0 mmol) was dissolved in 1 mL of DCM, and *m*-chloroperoxybenzoic acid (1 equiv) was added. The solution stirred at room temperature for 4 h. After completion of the reaction, the solvent was removed under reduced pressure and the residue was purified using flash chromatography (DCM:MeOH (9:1)).

4-(2-(4-Bromophenyl)acetyl)-1-thia-4,7,13-triazacyclopentadecane-6,12-dione 1-Oxide 7a. The product was obtained as a white solid (65%, 0.264 g, mp 205–207 °C); ^1H NMR (500 MHz, methanol- d_4) δ 7.36 (dd, $J = 17.7$, 8.0 Hz, 2H), 7.13 (d, $J = 8.1$ Hz, 1H), 7.06 (d, $J = 8.0$ Hz, 1H), 5.27–5.23 (m, 4H), 5.17 (s, 2H), 3.82 (s, 1H), 3.74 (s, 1H), 3.50 (d, $J = 8.8$ Hz, 2H), 3.23–3.14 (m, 5H), 2.18–2.05 (m, 2H), 1.59–1.33 (m, 4H). ^{13}C NMR (126 MHz, methanol- d_4) δ 173.1, 172.0, 168.6, 134.6, 133.2, 130.0, 126.6, 125.4, 118.8, 61.2, 54.5, 51.8, 51.7, 35.7, 35.4, 26.0, 22.0, 17.6. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_4\text{SBr}$ 472.0900; found 472.0901.

General Procedure and Analytical Data for the Synthesis of Sulfone Macrocycle. Macrocycle **6q** (1.0 mmol) was dissolved in 1 mL of DCM, and *m*-chloroperoxybenzoic acid (4 equiv) was added. The solution stirred at room temperature for 4 h. After completion of the reaction, the solvent was removed under reduced pressure and the residue was purified using flash chromatography (DCM:MeOH (9:1)).

4-(2-(4-Bromophenyl)acetyl)-1-thia-4,7,13-triazacyclopentadecane-6,12-dione 1,1-Dioxide 7b. The product was obtained as a white solid (77%, 0.375 g, mp 211–212 °C); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.81 (d, $J = 17.2$ Hz, 1H), 7.65 (s, 1H), 7.35 (dd, $J = 16.2$, 7.9 Hz, 2H), 7.14 (d, $J = 8.0$ Hz, 1H), 7.04 (d, $J = 8.0$ Hz, 1H), 3.99 (d, $J = 22.7$ Hz, 2H), 3.86–3.72 (m, 2H), 3.54 (d, $J = 14.6$ Hz, 3H), 3.38 (t, $J = 7.6$ Hz, 1H), 3.30 (d, $J = 7.7$ Hz, 1H), 3.27–3.09 (m, 5H), 2.57–2.51 (m, 1H), 2.12 (d, $J = 7.3$ Hz, 2H), 1.63–1.37 (m, 4H). ^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ 174.6, 172.5, 169.7, 140.1, 132.4, 132.1, 132.0, 125.0, 55.2, 54.9, 53.2, 52.9, 43.1, 40.6, 38.6, 36.0, 34.8, 29.4, 23.8. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_5\text{SBr}$ 488.0849; found 488.0848.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.7b02984.

NMR spectra, crystal structure determinations, and virtual library synthesis (PDF)

CIF data for **6c** (CIF)

CIF data for **6e** (CIF)

CIF data for **6l** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: a.s.s.dömling@rug.nl, www.drugdesign.nl.

ORCID

Eman M. M. Abdelraheem: 0000-0002-9008-2729

Justyna Kalinowska-Tłuścik: 0000-0001-7714-1651

Shabnam Shaabani: 0000-0001-5546-7140

Alexander Dömling: 0000-0002-9923-8873

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The work was financially supported by NIH 2R01GM097082-05, the Innovative Medicines Initiative (grant agreement no.115489), and also European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in-kind contribution. Funding came also from the European Union's Horizon 2020 research and innovation programme under MSC ITN "Accelerated Early Stage Drug Discovery" (no. 675555), CoFund ALERT (no. 665250). The work was supported by the European Regional Development Fund in the framework of the Polish Innovation Economy Operational Program (contract no. POIG.02.01.00-12-023/08). Eman M. M. Abdelraheem was supported by the Egyptian government.

■ REFERENCES

- (1) Domling, A.; Ugi, I. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168.
- (2) Domling, A. *Curr. Opin. Chem. Biol.* **2002**, *6*, 306.
- (3) Ruijter, E.; Scheffelaar, R.; Orru, R. V. A. *Angew. Chem., Int. Ed.* **2011**, *50*, 6234.
- (4) Domling, A.; Wang, W.; Wang, K. *Chem. Rev.* **2012**, *112*, 3083.
- (5) Koopmanschap, G.; Ruijter, E.; Orru, R. V. A. *Beilstein J. Org. Chem.* **2014**, *10*, 544.
- (6) Sinha, M.K.; Khoury, K.; Herdtweck, E.; Domling, A. *Org. Biomol. Chem.* **2013**, *11*, 4792.
- (7) Brauch, S.; van Berkel, S. S.; Westermann, B. *Chem. Soc. Rev.* **2013**, *42*, 4948.
- (8) Marsault, E.; Peterson, M. L. *J. Med. Chem.* **2011**, *54*, 1961.
- (9) Driggers, E. M.; Hale, S. P.; Lee, J.; Terrett, N. K. *Nat. Rev. Drug Discovery* **2008**, *7*, 608.
- (10) Failli, A.; Immer, H.; Gotz, M. *Can. J. Chem.* **1979**, *57*, 3257.
- (11) Wessjohann, L. A.; Voigt, B.; Rivera, D. G. *Angew. Chem., Int. Ed.* **2005**, *44*, 4785.
- (12) Hili, R.; Rai, V.; Yudin, A. K. *J. Am. Chem. Soc.* **2010**, *132*, 2889.
- (13) Jebail, M. J.; Ng, A. H. C.; Rai, V.; Hili, R.; Yudin, A. K.; Wheeler, A. R. *Angew. Chem., Int. Ed.* **2010**, *49*, 8625.
- (14) Liao, G. P.; Abdelraheem, E. M. M.; Neochoritis, C. G.; Kurpiewska, K.; Kalinowska-Tluscik, J.; McGowan, D. C.; Dömling, A. *Org. Lett.* **2015**, *17*, 4980.
- (15) Madhavachary, R.; Abdelraheem, E. M. M.; Rossetti, A.; Twarda-Clapa, A.; Musielak, B.; Kurpiewska, K.; Kalinowska-Tluscik, J.; Holak, T. A.; Dömling, A. *Angew. Chem., Int. Ed.* **2017**, *56*, 10725.
- (16) Abdelraheem, E. M. M.; de Haan, M. P.; Patil, P.; Kurpiewska, K.; Kalinowska-Tluscik, J.; Shaabani, S.; Domling, A. *Org. Lett.* **2017**, *19*, 5078.
- (17) Matsson, P.; Kihlberg, J. *J. Med. Chem.* **2017**, *60*, 1662.
- (18) Pye, C. R.; Hewitt, W. M.; Schwochert, J.; Haddad, T. D.; Townsend, C. E.; Etienne, L.; Lao, Y.; Limberakis, C.; Furukawa, A.; Mathiowetz, A. M.; Price, D. A.; Liras, S.; Lokey, R. S. *J. Med. Chem.* **2017**, *60*, 1665.